



PEPTECH BIOSCIENCES LIMITED

(Titan Biotech Ltd. Group)

G-1, 636, RIICO Industrial Area, Chopanki, Bhiwadi-301019 (Raj.)
CIN : U33110RJ2011PLC037007 | E-mail : sales1@peptechbio.com

Product Specification

AMINOFERT-MAGNESIUM

(MAGNESIUM AMINO ACID CHELATE - MG 6%)

AC 006

‘Aminofert-Magnesium’ Magnesium Amino Acid Chelated Powder is prepared for foliar application on plants to correct nutrient deficiency that may limit crop growth and yield. It is non-toxic to plants when applied. For best results, apply Aminofert-Magnesium soluble powder as per recommendations based on plant tissue or soil analysis. It is approved for organic production.

Aminofert-Magnesium (Magnesium Amino Acid Chelated Powder) contains 6% as Magnesium and 30% amino acids (as protein hydrolysed).

Aminofert-Magnesium (Magnesium Amino Acid Chelated Powder) is creamish to off white color free flowing powder, completely soluble in water and clear solution.

DEFICIENCY SYMPTOMS:

- Leaves become distorted, small, thick and brittle.
- Growing points start dying.
- Watery patches develop in storage tissues.
- Fruit development is irregular and malformed.
- Root growth is impaired.

STANDARD SPECIFICATION:

PHYSICAL PARAMETERS

S.L. No.	PRODUCT PARAMETERS	SPECIFICATION
1.	Appearance	Light creamish to white color free flowing powder.
2.	Solubility (10% soln.)	Completely soluble in water.

CHEMICAL PARAMETERS

S.L. No.	PRODUCT PARAMETERS	SPECIFICATION
1.	pH (2% soln. at 25°C)	5.0-7.5
2.	Magnesium Content (Mg)	NLT-6.0%
3.	Total Nitrogen Content	NLT-5.0%
4.	Amino acid Contents	NLT-30.0%
5.	Loss On Drying (at 105°C)	NMT-7.0%

BENEFITS:

- Magnesium Amino Acid Chelates helps the plant at the stage of growth and severity of deficiency.

DIRECTIONS:

Apply Aminofert-Magnesium by a spray method with an adequate amount of water that will provide complete coverage of the plants. Use of a non-ionic wetting agent may improve spray coverage of certain hard to wet plants.



PEPTECH BIOSCIENCES LIMITED

 (Titan Biotech Ltd. Group)

G-1, 636, RIICO Industrial Area, Chopanki, Bhiwadi-301019 (Raj.)
CIN : U33110RJ2011PLC037007 | E-mail : sales1@peptechbio.com

Product Specification

AMINOFERT-MAGNESIUM

(MAGNESIUM AMINO ACID CHELATE - MG 6%)

AC 006

Aminofert-Magnesium soluble powder can be included in a regular spray program on crops and compatible with other spray material.

The rate of application will depend on the crops, stage of growth, and severity of deficiency. The maximum recommended rates are for mature, full-sized plants, Reduce the rates proportionately when spraying smaller plants.

Aminofert-Magnesium soluble powder may be used on legumes, grain crops, root crops, cucurbits, cole crops, leafy vegetables, woody and herbaceous ornamentals, deciduous fruits, tropical and subtropical fruits, and many other crops.

DOSAGE:

Field Crops & Vegetables

Apply Aminofert-Magnesium 200 to 400 gm per acre during periods of rapid growth or nutritional stress. The application may be repeated twice or more times through the growth season.

Tree Crops

Make an application of Aminofert-Magnesium 400 to 800 gm per acre after the beginning of active growth. The application may be repeated at 2 to 4 weeks intervals through the growth season.

Grapes & Berries

Make an application of Aminofert-Magnesium 200 to 400 gm per acre after active growth begins. The application may be repeated at intervals of one week or more through the vegetative growth period.

INSTRUCTIONS:

Initially Soluble Powders will require special mixing attention. Like Soluble Powders have a tendency to clump when introduced to liquids. Because of this, take every step possible within the requirements to make these materials as user-friendly as possible.

Care should be taken to ensure that the materials are fully mixed and disposed throughout the spray tank. One way to assure proper dispersion is to place the required amount of powder into a bucket of water. After fully mixing, the product be added to the spray tank. You can then mix the next spray tank worth of powder in a bucket and let it hydrate while spraying the first load. This makes the mixing much easier.

Other methods of dispersal:

- Use an inductor system.
- Wash the product through the anti-splash screen on the top of the spray tank.

STD. Packing : 25 Kg Paper Bag.

Best Before : 36 Months.

Storage : Store the material below 25°C, protect from direct sunlight.